

What can we learn from $B \rightarrow a_1(1260)(b_1(1235))\pi(K)$ decays?

Wei Wang^a, Run-Hui Li^{b,a} and Cai-Dian Lü^{a,c}

^a *Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, P.R. China*

^b *Physics Department, Shandong University, Jinan 250100, P.R. China*

^c *Theoretical Physics Center for Science Facilities, CAS, Beijing 100049, P.R. China*

We investigate the $B \rightarrow a_1(1260)(b_1(1235))\pi(K)$ decays under the factorization scheme and find many discrepancies between theoretical predictions and the experimental data. In the tree dominated processes, large contributions from color-suppressed tree diagrams are required in order to accommodate with the large decay rates of $B^- \rightarrow a_1^0 \pi^-$ and $B^- \rightarrow a_1^- \pi^0$. For $\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-$ decays which are both induced by $b \rightarrow s$ transition, theoretical predictions on their decay rates are larger than the data by a factor of 2.8 and 5.5, respectively. Large electro-weak penguins or some new mechanism are expected to explain the branching ratios of $B^- \rightarrow b_1^0 K^-$ and $B^- \rightarrow a_1^- \bar{K}^0$. The soft-collinear-effective-theory has the potential to explain large decay rates of $B^- \rightarrow a_1^0 \pi^-$ and $B^- \rightarrow a_1^- \pi^0$ via a large hard-scattering form factor $\zeta_J^{B \rightarrow a_1}$. We will also show that, with proper charming penguins, predictions on the branching ratios of $\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-$ can also be consistent with the data.

PACS numbers: 13.25.Hw, 14.40.Cs

I. INTRODUCTION

Since the first measurement on $B^0/\bar{B}^0 \rightarrow a_1^\mp(1260)\pi^\pm$ decays reported by BaBar and Belle collaborations [1, 2, 3], many charmless B decays into a pseudo-scalar and an axial-vector meson have been observed. Among the 18 $B \rightarrow a_1(1260)(b_1(1235))\pi(K)$ ¹ decay channels, 10 of them have been measured with large branching ratios. Besides decay rates, direct CP asymmetries in some $B \rightarrow (a_1, b_1)K$ channels and time-dependent CP asymmetries in $B^0/\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$ and $B^0/\bar{B}^0 \rightarrow b_1^\pm \pi^\mp$ were also studied in the two B factories [4, 5, 6, 7, 8, 9]. Without any doubt, these results are helpful to investigate production mechanisms of axial-vectors in B decays, extract hadronic parameters such as strong phases in $B \rightarrow AP$ decays and probe the structures of axial-vectors.

Charmless two-body $B \rightarrow AP$ decays have received considerable theoretical efforts [10, 11, 12, 13, 14]. Among these predictions, many of them are not consistent with each other:

¹ In the following, we will use $a_1(b_1)$ to denote the $a_1(1260)(b_1(1235))$ meson for simplicity.

most predictions by Calderon, Munoz and Vera [12] are larger than predictions given by Laporta, Nardulli and Pham [11] and the QCD factorization (QCDF) approach. Predictions on $B \rightarrow a_1\pi$ by Laporta, Nardulli and Pham (using the second sets of form factors) are very close to results in the QCDF approach. However there are large discrepancies in other predictions (See Ref. [14] for a detailed comparison between these theoretical predictions). Many results of the QCDF approach agree with the experimental data, but there still exist some deviations.

In the present paper, we intend to analyze the 18 $B \rightarrow AP$ decays with the help of experimental data. We try to check whether these problems can be removed in the perturbative QCD (PQCD) approach and the soft-collinear-effective-theory (SCET). Another objective is to extract the $B \rightarrow A$ form factors through $\bar{B}^0 \rightarrow a_1^+\pi^-$ and $\bar{B}^0 \rightarrow b_1^\pm\pi^\mp$ decays.

II. NAIVE FACTORIZATION APPROACH

The effective Hamiltonian describing $b \rightarrow D(D = d, s)$ transitions are given by [15]:

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \left\{ \sum_{q=u,c} V_{qb}V_{qD}^* \left[C_1 O_1^q + C_2 O_2^q + \sum_{i=3}^{10} C_i O_i \right] \right\} + \text{H.c.}, \quad (1)$$

where $V_{qb(D)}$ are the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. Functions O_i are the local four-quark operators, while functions C_i are the corresponding Wilson coefficients. It is convenient to define combinations a_i of the Wilson coefficients:

$$\begin{aligned} a_1 &= C_2 + C_1/3, \quad a_2 = C_1 + C_2/3, \\ a_i &= C_i + C_{i+1}/N_c \quad (i = 3, 5, 7, 9), \\ a_i &= C_i + C_{i-1}/N_c \quad (i = 4, 6, 8, 10). \end{aligned} \quad (2)$$

There exist a hierarchy for the Wilson coefficients:

$$a_1 \gg \max[a_2, a_{3-10}]. \quad (3)$$

For tree-dominated processes $B^0/\bar{B}^0 \rightarrow a_1^\pm\pi^\mp$, the factorization formulae can be written as:

$$\begin{aligned} \mathcal{A}(\bar{B}^0 \rightarrow a_1^+\pi^-) &= \frac{G_F}{\sqrt{2}} m_B^2 f_\pi V_0^{B \rightarrow a_1} \{ V_{ub}V_{ud}^* [a_1 + a_4 + a_{10} + r_\pi(a_6 + a_8)] \\ &\quad + V_{cb}V_{cd}^* [a_4 + a_{10} + r_\pi(a_6 + a_8)] \}, \end{aligned} \quad (4)$$

$$\mathcal{A}(\bar{B}^0 \rightarrow \pi^+a_1^-) = \frac{G_F}{\sqrt{2}} m_B^2 f_{a_1} f_+^{B \rightarrow \pi} \{ V_{ub}V_{ud}^* [a_1 + a_4 + a_{10}] + V_{cb}V_{cd}^* [a_4 + a_{10}] \}, \quad (5)$$

where $r_\pi = 2m_0^\pi/m_B$ with m_0^π the chiral scale parameter for pion. The CKM matrix elements for tree operators $|V_{ub}V_{ud}^*| \sim 4 \times 10^{-3}$ have the same order magnitude with those for penguin

operators $|V_{cb}V_{cd}^*| \sim 8 \times 10^{-3}$. Because of the hierarchy in the Wilson coefficients, penguin contributions from the operators O_{3-10} are small compared with those from tree operators. Thus penguin contributions can be neglected in the study of branching ratios (but crucial to CP asymmetries). Combined with the $\bar{B}^0 \rightarrow \pi^+\pi^-$ data [16]

$$\mathcal{BR}(\bar{B}^0 \rightarrow \pi^+\pi^-) = (5.16 \pm 0.22) \times 10^{-6}, \quad (6)$$

we arrive at the a_1 meson decay constant and $B \rightarrow a_1$ form factor:

$$f_{a_1} = [2.02 \pm 0.26 \pm 0.04 + \mathcal{O}(\frac{a_{3-10}}{a_1})]f_\pi, \quad V_0^{B \rightarrow a_1} = (1.55 \pm 0.28 \pm 0.03 + \mathcal{O}(\frac{a_{3-10}}{a_1}))f_+^{B \rightarrow \pi} \quad (7)$$

where the uncertainties are from the experimental results for branching ratios. As a rough estimation, we take $f_\pi = 131$ MeV and $f_+^{B \rightarrow \pi} = 0.25$ which corresponds to $f_{a_1} = (265 \pm 34 \pm 6)$ MeV and $V_0^{B \rightarrow a_1} = 0.39 \pm 0.07 \pm 0.01$. These results are well consistent with predictions based on the PQCD approaches [17] and light-cone sum rules [18, 19].

Now we come to the two channels $B^- \rightarrow a_1^0\pi^-$ and $B^- \rightarrow a_1^-\pi^0$ whose factorization formulae are given by:

$$\begin{aligned} \sqrt{2}\mathcal{A}(B^- \rightarrow \pi^- a_1^0) &= \frac{G_F}{\sqrt{2}}m_B^2 f_\pi V_0^{B \rightarrow a_1} \{V_{ub}V_{ud}^*[a_1 + a_4 + a_{10} + r_\pi(a_6 + a_8)] \\ &\quad + V_{cb}V_{cd}^*[a_4 + a_{10} + r_\pi(a_6 + a_8)]\} \\ &\quad + \frac{G_F}{\sqrt{2}}m_B^2 f_{a_1} f_+^{B \rightarrow \pi} \left\{ V_{ub}V_{ud}^*[a_2 - a_4 + \frac{1}{2}a_{10}] + V_{cb}V_{cd}^*[-a_4 + \frac{1}{2}a_{10}] \right\}, \quad (8) \\ \sqrt{2}\mathcal{A}(B^- \rightarrow \pi^0 a_1^-) &= \frac{G_F}{\sqrt{2}}m_B^2 f_\pi V_0^{B \rightarrow a_1} \left\{ V_{ub}V_{ud}^*[a_2 - a_4 + \frac{1}{2}a_{10} + r_\pi(-a_6 + \frac{1}{2}a_8)] \right. \\ &\quad \left. + V_{cb}V_{cd}^*[-a_4 + \frac{1}{2}a_{10} + r_\pi(-a_6 + \frac{1}{2}a_8)] \right\} \\ &\quad + \frac{G_F}{\sqrt{2}}m_B^2 f_{a_1} f_+^{B \rightarrow \pi} \{V_{ub}V_{ud}^*[a_1 + a_4 + a_{10}] + V_{cb}V_{cd}^*[a_4 + a_{10}]\}. \quad (9) \end{aligned}$$

Because of the small values of a_{3-10} , the penguin contributions can be safely neglected:

$$\sqrt{2}\mathcal{A}(B^- \rightarrow \pi^- a_1^0) = \frac{G_F}{\sqrt{2}}m_B^2 V_{ub}V_{ud}^*[a_1 f_\pi V_0^{B \rightarrow a_1} + a_2 f_{a_1} f_+^{B \rightarrow \pi}], \quad (10)$$

$$\sqrt{2}\mathcal{A}(B^- \rightarrow \pi^0 a_1^-) = \frac{G_F}{\sqrt{2}}m_B^2 V_{ub}V_{ud}^*[a_2 f_\pi V_0^{B \rightarrow a_1} + a_1 f_{a_1} f_+^{B \rightarrow \pi}]. \quad (11)$$

Furthermore, in the hierarchy of $a_2 \ll a_1$, branching ratios are required to satisfy the following relation:

$$\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+\pi^-) = 2\mathcal{BR}(B^- \rightarrow \pi^- a_1^0), \quad \mathcal{BR}(\bar{B}^0 \rightarrow \pi^+ a_1^-) = 2\mathcal{BR}(B^- \rightarrow a_1^-\pi^0). \quad (12)$$

But the experimental data shows:

$$\mathcal{BR}(B^- \rightarrow \pi^0 a_1^-) > \mathcal{BR}(\bar{B}^0 \rightarrow a_1^-\pi^+), \quad \mathcal{BR}(B^- \rightarrow \pi^- a_1^0) > \mathcal{BR}(\bar{B}^0 \rightarrow a_1^+\pi^-), \quad (13)$$

which is dramatically different. This situation is very similar with that in $B \rightarrow \pi\pi$ decays: the branching ratio of $B^- \rightarrow \pi^0\pi^-$ is measured with almost equal magnitude with $\mathcal{BR}(\bar{B}^0 \rightarrow \pi^-\pi^+)$ but it is expected as one half of $\mathcal{BR}(\bar{B}^0 \rightarrow \pi^-\pi^+)$. To solve these problems, an efficient way is to enhance the color-suppressed contribution which is proportional to a_2 . For example, if the Wilson coefficient a_2 can be enhanced to 0.5, the branching ratios of $\mathcal{BR}(B^- \rightarrow \pi^0 a_1^-)$ and $\mathcal{BR}(B^- \rightarrow \pi^- a_1^0)$ are predicted as 20.0×10^{-6} and 16.7×10^{-6} , where we have utilized the experimental data on branching ratios of $B^0/\bar{B}^0 \rightarrow \pi^\pm a_1^\mp$. And these results are well consistent with the experimental data.

The decay constant of b_1 vanishes because of the G-parity, thus $\bar{B}^0 \rightarrow \pi^+ b_1^-$ is factorization-suppressed and only the $\bar{B}^0 \rightarrow \pi^- b_1^+$ decay survives. From the experimental results collected in table I, we can infer that the form factors of $B \rightarrow a_1$ and $B \rightarrow b_1$ are almost equal in magnitude at maximally recoiling: $|V_0^{B \rightarrow a_1}(q^2 = 0)| \simeq |V_0^{B \rightarrow b_1}(q^2 = 0)| \simeq 0.35$. One should be careful that the two form factors have different signs, if we use LCDAs of a_1 and b_1 evaluated by the QCD sum rules. The absolute value of these form factors can be checked by the future measurements on semi-leptonic $B \rightarrow A$ decays such as $\bar{B}^0 \rightarrow (a_1^+, b_1^+) l^- \bar{\nu}$.

Flavor structures of $\bar{B}^0 \rightarrow b_1^+ K^-$ and $\bar{B}^0 \rightarrow a_1^+ K^-$ are the same with each other, thus they have the same factorization formulae:

$$\begin{aligned} \mathcal{A}(\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-) = & \frac{G_F}{\sqrt{2}} m_B^2 f_K V_0^{B \rightarrow (a_1, b_1)} \{ V_{ub} V_{us}^* [a_1 + a_4 + a_{10} + r_K(a_6 + a_8)] \\ & + V_{cb} V_{cs}^* [a_4 + a_{10} + r_K(a_6 + a_8)] \}. \end{aligned} \quad (14)$$

The same Wilson coefficients and almost equal form factors will induce almost equal branching ratios for $\bar{B}^0 \rightarrow a_1^+ K^-$ and $\bar{B}^0 \rightarrow b_1^+ K^-$. To reduce the uncertainties, we will utilize the $\bar{B}^0 \rightarrow \pi^+ K^-$ decay which also has the same flavor structures with $\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-$. The only difference between the three modes is the different form factors which can be extracted from tree-dominated processes $\bar{B}^0 \rightarrow \pi^+ \pi^-$ and $\bar{B}^0 \rightarrow (a_1^+, b_1^+) \pi^-$ decays. The branching ratio of $\bar{B}^0 \rightarrow \pi^+ K^-$ has been measured as [16]:

$$\mathcal{BR}(\bar{B}^0 \rightarrow \pi^+ K^-) = (19.4 \pm 0.6) \times 10^{-6}, \quad (15)$$

which implies:

$$\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+ K^-) = 45.9 \times 10^{-6}, \quad \mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ K^-) = 41.0 \times 10^{-6}. \quad (16)$$

Comparing with the experimental measurements in table I, we see that our theoretical prediction on $\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+ K^-)$ is 2.8 times larger while the prediction on $\mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ K^-)$ is 5.5 times larger. This discrepancy should be clarified by the theoretical studies with next-to-leading order corrections and improved experimental measurements.

Besides $\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-$ decays, $B^- \rightarrow a_1^- \bar{K}^0$ and $B^- \rightarrow b_1^0 K^-$ decays are also measured by experimentalists whose factorization formulae are:

$$\begin{aligned} \mathcal{A}(B^- \rightarrow a_1^- \bar{K}^0) = & \frac{G_F}{\sqrt{2}} m_B^2 f_K V_0^{B \rightarrow a_1} \left\{ V_{ub} V_{us}^* \left[a_4 - \frac{1}{2} a_{10} + r_K \left(a_6 - \frac{1}{2} a_8 \right) \right] \right. \\ & \left. + V_{cb} V_{cs}^* \left[a_4 - \frac{1}{2} a_{10} + r_K \left(a_6 - \frac{1}{2} a_8 \right) \right] \right\}, \end{aligned} \quad (17)$$

$$\begin{aligned} \sqrt{2} \mathcal{A}(B^- \rightarrow b_1^0 K^-) = & \frac{G_F}{\sqrt{2}} m_B^2 f_K V_0^{B \rightarrow b_1} \left\{ V_{ub} V_{us}^* [a_1 + a_4 + a_{10} + r_K (a_6 + a_8)] \right. \\ & \left. + V_{cb} V_{cs}^* [a_4 + a_{10} + r_K (a_6 + a_8)] \right\}. \end{aligned} \quad (18)$$

In these $b \rightarrow s$ transitions, the CKM matrix elements for penguin operators are $|V_{cb} V_{cs}^*| \sim 40 \times 10^{-3}$ and those for tree operators are $|V_{ub} V_{us}^*| \sim 0.8 \times 10^{-3}$. Recalling the values for the Wilson coefficient combinations: $a_1 \sim 1$ and $a_4 \sim a_6 \sim -0.03$, we can see that contributions from tree operators with the coefficient a_1 are smaller than that from penguin operators at least by a factor of 2 in magnitude. In order to characterize the contribution from tree operators and symmetry breaking effects between B^- and $\bar{B}^0$ mesons, it is useful to define the two ratios:

$$R_1 \equiv \frac{\mathcal{BR}(B^- \rightarrow a_1^- \bar{K}^0)}{\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+ K^-)} \times \frac{\tau_{\bar{B}^0}}{\tau_B}, \quad R_2 \equiv \frac{\mathcal{BR}(B^- \rightarrow b_1^0 K^-)}{\mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ K^-)} \times \frac{\tau_{\bar{B}^0}}{\tau_B}, \quad (19)$$

where τ is the lifetime of B meson. Neglecting tree operators and electro-weak penguins, the ratios obey the limit:

$$R_1 = 1, \quad R_2 = 0.5, \quad (20)$$

which are quite different from the experimental results:

$$R_1^{\text{exp.}} = 2.00 \pm 0.59, \quad R_2^{\text{exp.}} = 1.15 \pm 0.34. \quad (21)$$

The difference between the two channels in the ratio R_1 is the tree operator and electroweak penguin operators. Since the contribution of tree operator is smaller than QCD penguins and the two kinds of amplitudes are perpendicular with each other due to the CKM angle γ close to 90° , the tree operator can not change the branching ratio of $\bar{B}^0 \rightarrow a_1^+ K^-$ too much. Thus this does not improve theoretical predictions on R_1 . Large electro-weak penguins may help us to diminish the large deviation for R_1 . In the $\bar{B}^0 \rightarrow b_1^+ K^-$ and $B^- \rightarrow b_1^0 K^-$ decays, the factorization formulae are exactly the same since the b_1 decay constant vanishes. Thus in order to explain the large ratio R_2 , one needs some mechanism beyond factorization to enhance the ratio of R_2 by roughly 2.5.

In the above, we have analyzed the charmless non-leptonic $B \rightarrow AP$ data under the factorization approach. The decay constant of a_1 meson and $B \rightarrow a_1, b_1$ form factors V_0 are

extracted from the $\bar{B}^0 \rightarrow a_1\pi$ and $\bar{B}^0 \rightarrow b_1\pi$ decays. The form factors are consistent with the predictions evaluated in light-cone-sum-rules and the PQCD approach. But there exist several problems which can be summarized as:

- The Wilson coefficient combination a_2 needs to be enhanced to $a_2 = 0.5$ in order to solve the problem in $B^- \rightarrow a_1^- \pi^0$ and $B^- \rightarrow a_1^0 \pi^-$.
- Since the form factor $B \rightarrow a_1$ and $B \rightarrow b_1$ are almost equal in magnitude, the $\bar{B}^0 \rightarrow a_1^+ K^-$ and $\bar{B}^0 \rightarrow b_1^+ K^-$ decays should possess similar and large branching ratios. Compared with the experimental data, theoretical predictions need to be reduced by the factors of 2.8 and 5.5, respectively.
- $B^- \rightarrow a_1^- \bar{K}^0$ and $B^- \rightarrow b_1^0 K^-$ are related to $\bar{B}^0 \rightarrow (a_1^+, b_1^+) K^-$ through relations given in Eq. (19) which also have large deviations from the data.

III. THE SOFT-COLLINEAR EFFECTIVE THEORY

The recent development of SCET makes the analysis of $B \rightarrow M_1 M_2$ decays on a more rigorous foundation. The SCET is a powerful method to systematically separate the dynamics at different scales: hard scale m_b (b quark mass), hard intermediate scale $\mu_{hc} = \sqrt{m_b \Lambda_{QCD}}$, soft scale and to sum large logs using the renormalization group technics. Integrating out the hard fluctuations, we arrive at the intermediate effective theory- SCET_I where the factorization formulae for $B \rightarrow M_1 M_2$ decays to leading power in $\lambda \equiv \sqrt{\Lambda_{QCD}/m_b}$ are given by:

$$\begin{aligned} \mathcal{A}(B \rightarrow M_1 M_2) = & \frac{G_F}{\sqrt{2}} m_B^2 \left\{ f_{M_1} \int du \phi_{M_1}(u) T_1(u) \zeta^{B \rightarrow M_2} \right. \\ & \left. + f_{M_1} \int du \phi_{M_1}(u) \int dz T_{1J}(u, z) \zeta_J^{B \rightarrow M_2}(z) + (1 \leftrightarrow 2) \right\}, \quad (22) \end{aligned}$$

where functions ζ and ζ_J also enter into the heavy-to-light form factors. $T_1(u)$ and $T_{1J}(u, z)$ are hard kernels which can be calculated using perturbation theory. With the hard-collinear fluctuation integrated out, the final effective theory-SCET_{II} is obtained where the function ζ_J can be factorized into convolutions of LCDAs with hard kernels:

$$\zeta_J^{B \rightarrow M_2}(z) = \phi_B(\omega) \otimes J(z, \omega, v) \otimes \phi_{M_2}(v). \quad (23)$$

$J(z, \omega, v)$ is the hard kernel and ϕ_B and ϕ_{M_1, M_2} are the light-cone distribution amplitudes (LCDAs). With our knowledge on these LCDAs, one can predict the decay amplitude by convoluting the LCDAs with the perturbatively calculated hard kernels. But there is another alternative way for phenomenological studies: one can fit experimental results, including

branching ratios and CP asymmetries, to determine essential non-perturbative inputs. Note that in this way, no expansions in $\alpha_s(\sqrt{m_b \Lambda_{QCD}})$ are needed and thus the exploration of the convergence is spontaneously avoided. This method is especially useful at tree level: $T_1(u)$ is a constant and $T_{1J}(u, z)$ is a function of one argument u . It leads to a rather simple form for decay amplitudes:

$$\mathcal{A}(B \rightarrow M_1 M_2) = \frac{G_F}{\sqrt{2}} m_B^2 \left\{ f_{M_1} T_1 \zeta^{B \rightarrow M_2} + f_{M_1} \int du \phi_{M_1}(u) T_{1J}(u) \zeta_J^{B \rightarrow M_2} + (1 \leftrightarrow 2) \right\} \quad (24)$$

where the functions $\zeta^{B \rightarrow M_2}$ and

$$\zeta_J^{B \rightarrow M_2} = \int dz \zeta_J^{B \rightarrow M_2}(z) \quad (25)$$

are treated as non-perturbative parameters to be fitted from the data. With the help of the flavor SU(3) symmetry, the $B \rightarrow AP$ decays involve only 6 parameters:

$$\zeta^{B \rightarrow P}, \quad \zeta_J^{B \rightarrow P}, \quad \zeta^{B \rightarrow ^1P_1}, \quad \zeta_J^{B \rightarrow ^1P_1}, \quad \zeta^{B \rightarrow ^3P_1}, \quad \zeta_J^{B \rightarrow ^3P_1}, \quad (26)$$

which contribute to the $B \rightarrow P$ and $B \rightarrow A$ form factors.

Including the non-perturbative contributions from loop diagrams involving $c\bar{c}$ [20, 21, 22, 23, 24, 25], the SCET can successfully explain most of $B \rightarrow PP$ and $B \rightarrow VP$ decays [21, 26, 27]. This phenomenological approach has many important features. In $b \rightarrow d$ transitions such as $\bar{B}^0 \rightarrow \pi^+ \pi^-$, tree operators provide the dominant contributions and contributions from charming penguins and penguin operators are sub-leading. From the experience in $B \rightarrow PP$ and $B \rightarrow VP$ phenomenological study, we know that the hard-scattering form factor ζ_J is potentially large. Furthermore, as we have shown in Ref. [27], the corresponding Wilson coefficient is of order 1 which amounts to a large effective Wilson coefficient a_2 . Here we take $\bar{B}^0 \rightarrow a_1^- \pi^+$ and $B^- \rightarrow a_1^- \pi^0$ as an example: if hard-scattering form factors are equal with soft form factors for pion and a_1 meson: $\zeta = \zeta_J$, the effective Wilson coefficient equals to $a_2 \simeq \frac{\zeta_J}{\zeta + \zeta_J} = 0.5$. Thus it is easy to solve the problems in $B \rightarrow a_1 \pi$ decays under the SCET framework.

For decays induced by $b \rightarrow s$ transition, since tree operators are suppressed by the CKM matrix elements $|V_{ub} V_{us}^* / (V_{cb} V_{cs}^*)| \sim 0.02$ and penguin operators have smaller Wilson coefficients ($\max[C_{3-10}] \ll \alpha_s(2m_c) C_1$), charming penguins play a significant role. Due to the non-perturbative nature, charming penguins are totally unknown from perturbation theory and needs to be extracted from data. This stuff depends on the three involved mesons: B meson, recoiling meson and emitted meson. Thus in order to predict physical observables, too many parameters for charming penguins are required. An efficient way to reduce the independent inputs is to utilize the flavor SU(3) symmetry and as a result only 8 parameters for charming penguins in $B \rightarrow AP$ decays are left. But even so, due to the lack of data, one

can always obtain proper branching ratios of $\bar{B}^0 \rightarrow b_1^+ K^-$ and $\bar{B}^0 \rightarrow a_1^+ K^-$ by adjusting charming penguins. Despite of that, there is another deficit: since the inputs, form factors and charming penguins, have been assumed to respect the SU(3) symmetry, large deviations of the ratios shown in Eqs. (20) and (21) can not be eliminated by the SCET either.

IV. THE PERTURBATIVE QCD APPROACH

There is another commonly-accepted approach to handle hadronic B decays: the perturbative QCD approach [28, 29, 30]. The basic idea of the PQCD approach is that it takes into account the transverse momentum of the valence quarks in hadrons. Decay amplitudes and form factors can be written as convolutions of wave functions with perturbatively hard kernels integrated over the longitudinal and transverse component. When considering radiative corrections, one encounters double logarithm divergences when soft and collinear momenta overlap. These large double logarithm can be resummed into the Sudakov factor. Loop corrections to the weak decay vertex also give rise to double logarithms in the threshold region. Resummation of this type of double logarithms leads to the Sudakov factor S_t . This factor decreases faster than any power of x as $x \rightarrow 0$ and changes the behavior at the end-point region. The Sudakov factor and threshold resummation make the PQCD approach more self-consistent. This approach have successfully explained the $B \rightarrow \pi\pi$ and $B \rightarrow \pi K$ decay rates and CP asymmetries [31] together with the proper polarizations in $B \rightarrow VV$ decays [32].

In the PQCD approach, the predicted $B \rightarrow a_1$ form factor [17] is consistent with the one derived from the data, thus our PQCD prediction on $\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+ \pi^-)$ is in good agreement with the data. But due to the small value of a_2 , the color-suppressed contribution is too small to explain the large decay rates of $B^- \rightarrow a_1^- \pi^0$ and $B^- \rightarrow a_1^0 \pi^-$. The investigations of next-to-leading order corrections in Ref. [33] show that the branching ratio of $B^- \rightarrow \pi^- \pi^0$ is enhanced by the factor 4.0/3.5 while $\bar{B}^0 \rightarrow \pi^+ \pi^-$ is reduced by 6.5/7.0. But even if we assume the same k factor for $B \rightarrow a_1 \pi$ decays, the PQCD predictions on $B^- \rightarrow a_1^- \pi^0$ and $B^- \rightarrow a_1^0 \pi^-$ are still smaller than the data. The PQCD prediction on the $B \rightarrow b_1$ form factor is large, thus the branching ratio of $\bar{B}^0 \rightarrow b_1^+ \pi^-$ is 2 times larger than the experimental data and the QCDF results. From the factorization formulae of $B \rightarrow PP$ decays given in the literature [30], one can see that the contributions from hard spectator scattering diagrams are small due to the cancelation between two diagrams where a gluon is attached to either the positive quark or the anti-quark in the emitted hadron. But if the emitted meson is a P -wave meson and the twist-2 LCDA is anti-symmetric (like a scalar or an axial-vector meson with quantum number $^{2S+1}L_J = ^1P_1$), the two diagrams give constructive contributions to make them sizable. For example, the large hard spectator scattering contributions to

$B^- \rightarrow b_1^0 \pi^-$ make $\mathcal{BR}(B^- \rightarrow b_1^0 \pi^-) > \frac{1}{2} \mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ \pi^-)$. Moreover, annihilation diagrams play an important role in the PQCD approach which often enters into decay amplitudes as imaginary. It provides the dominant strong phase which are essential to explain the large CP asymmetries. Thus unlike the situation in the QCDF approach, annihilation diagrams do not cancel with emission diagrams in $\bar{B}^0 \rightarrow b_1^+ K^-$ which results in much larger predictions on branching ratios of $\bar{B}^0 \rightarrow b_1^+ K^-$. Similar as the factorization approach, there are large differences between the PQCD approach predictions on ratios $R_{1,2}$ and those extracted from the data.

V. NUMERICAL RESULTS

In the PQCD framework and SCET framework, we calculate the decay rates, direct CP asymmetries and time-dependent CP asymmetries shown in table I,II,III and IV. We have adopted the same conventions with Ref. [14] for observables in time-dependent decay widths of $B \rightarrow a_1^\pm \pi^\mp$ and $B \rightarrow b_1^\pm \pi^\mp$ ². In the SCET calculation, we use the following values for the 14 inputs:

$$\begin{aligned}
\zeta^{B \rightarrow \pi} &= 0.12, & \zeta_J^{B \rightarrow \pi} &= 0.12, \\
\zeta^{B \rightarrow a_1} &= 0.17, & \zeta_J^{B \rightarrow a_1} &= 0.17, \\
\zeta^{B \rightarrow b_1} &= -0.16, & \zeta_J^{B \rightarrow b_1} &= -0.16, \\
|A_{cc}^{3P_1 P}| &= 40 \times 10^{-4}, & \arg[A_{cc}^{3P_1 P}] &= 160^\circ, \\
|A_{cc}^{P^3 P_1}| &= 40 \times 10^{-4}, & \arg[A_{cc}^{P^3 P_1}] &= 145^\circ, \\
|A_{cc}^{1P_1 P}| &= 40 \times 10^{-4}, & \arg[A_{cc}^{1P_1 P}] &= 155^\circ, \\
|A_{cc}^{P^1 P_1}| &= 35 \times 10^{-4}, & \arg[A_{cc}^{P^1 P_1}] &= 100^\circ.
\end{aligned} \tag{27}$$

We should point out that this set of inputs is presented by hand instead of any reasonable way. To test the sensitivities on these parameters, we show the first uncertainty in numerical results by varying the form factors by 0.03, 20% for magnitudes of charming penguins and 20° for the phases. The second uncertainty is from CKM matrix elements. In the PQCD calculation, we have used the same inputs as those in Ref. [17, 36, 37]. The theoretical uncertainties are from: (i) the hadronic inputs: decay constants of B meson, and shape parameters in the wave function of B meson; (ii) Λ_{QCD} , the hard scale t and the threshold resummation parameter c ; (iii) the CKM matrix elements V_{ub} and γ angle. The factorization formula for each type of diagrams in $B \rightarrow AP$ decays are the same with those in $B \rightarrow PP$

² The $\mathcal{A}_{a_1^\pm \pi^\mp}$ in the present paper correspond to $\mathcal{A}_{a_1^\mp \pi^\pm}$ defined in Ref. [16, 34, 35].

TABLE I: Theoretical predictions and experimental results [1, 2, 3, 4, 5, 6, 7, 8, 9] on branching ratios (in unit of 10^{-6}) of $B \rightarrow a_1(b_1)\pi(K)$ decays. The QCDF predictions are quoted from Ref. [14]. In the PQCD approach, the uncertainties are from: (i) the hadronic inputs: decay constants of B meson, and shape parameters in the wave function of B meson; (ii) Λ_{QCD} , the hard scale t and the threshold resummation parameter c ; (iii) the CKM matrix elements V_{ub} and γ angle. In the SCET framework, the uncertainties are from: (i) hadronic parameters: form factors and charming penguins; (ii) the CKM matrix elements.

channel	QCDF	PQCD	SCET	Exp.
$B^- \rightarrow a_1^- \pi^0$	$14.4^{+1.4+3.5+2.1}_{-1.3-3.2-1.9}$	$8.1^{+4.1+2.1+0.7}_{-2.7-1.2-0.9}$	$19.0^{+5.1+1.8}_{-4.7-1.7}$	$26.4 \pm 5.4 \pm 4.1$
$B^- \rightarrow a_1^0 \pi^-$	$7.6^{+0.3+1.7+1.4}_{-0.3-1.3-1.0}$	$6.7^{+2.9+2.8+0.5}_{-2.2-1.7-0.7}$	$17.2^{+4.7+1.7}_{-4.3-1.6}$	$20.4 \pm 4.7 \pm 3.4$
$\bar{B}^0 \rightarrow a_1^- \pi^+$	$23.4^{+2.3+6.2+1.9}_{-2.2-5.5-1.3}$	$15.7^{+8.3+5.9+1.2}_{-5.6-3.6-1.7}$	$17.0^{+5.8+1.6}_{-5.2-1.4}$	21.0 ± 5.4
$\bar{B}^0 \rightarrow a_1^+ \pi^-$	$9.1^{+0.2+2.2+1.7}_{-0.2-1.8-1.1}$	$12.7^{+5.6+6.2+0.9}_{-4.4-3.8-1.3}$	$10.7^{+2.5+1.0}_{-2.4-0.9}$	12.2 ± 4.5
$B^0/\bar{B}^0 \rightarrow a_1^+ \pi^-$	—	$28.1^{+13.8+12.0+2.1}_{-9.9-7.3-3.0}$	$28.2^{+6.5+2.6}_{-5.9-2.4}$	
$B^0/\bar{B}^0 \rightarrow a_1^- \pi^+$	—	$28.6^{+13.9+12.1+2.2}_{-10.1-7.4-3.0}$	$27.1^{+6.9+2.5}_{-6.2-2.3}$	
$\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$	$32.5^{+2.5+8.4+3.6}_{-2.4-7.3-2.4}$	$28.3^{+13.9+12.0+2.2}_{-10.0-7.4-3.0}$	$27.7^{+6.3+2.5}_{-5.7-2.3}$	31.7 ± 3.7
$\bar{B}^0 \rightarrow a_1^0 \pi^0$	$0.9^{+0.1+0.3+0.7}_{-0.1-0.2-0.3}$	$0.12^{+0.07+0.02+0.02}_{-0.04-0.03-0.02}$	$5.5^{+1.7+0.6}_{-1.5-0.6}$	
$B^- \rightarrow a_1^0 K^-$	$13.9^{+0.9+9.5+12.9}_{-0.9-5.1-4.9}$	$15.4^{+7.8+10.1+2.4}_{-5.4-5.5-2.5}$	$10.5^{+3.3+1.8}_{-2.9-1.5}$	
$B^- \rightarrow a_1^- \bar{K}^0$	$21.6^{+1.2+16.5+23.6}_{-1.1-8.5-11.9}$	$25.5^{+12.9+18.0+3.7}_{-9.2-10.2-3.9}$	$15.5^{+5.8+2.5}_{-5.0-2.1}$	$34.9 \pm 5.0 \pm 4.4$
$\bar{B}^0 \rightarrow a_1^+ K^-$	$18.3^{+1.0+14.2+21.1}_{-1.0-7.2-7.5}$	$20.6^{+10.2+14.6+3.2}_{-7.3-8.5-3.3}$	$15.8^{+5.6+2.7}_{-4.9-2.3}$	$16.3 \pm 2.9 \pm 2.3$
$\bar{B}^0 \rightarrow a_1^0 \bar{K}^0$	$6.9^{+0.3+6.1+9.5}_{-0.3-2.9-3.2}$	$8.0^{+3.9+6.4+1.2}_{-2.8-3.4-1.2}$	$6.3^{+2.5+1.0}_{-2.1-0.8}$	
$B^- \rightarrow b_1^- \pi^0$	$0.4^{+0.0+0.2+0.4}_{-0.0-0.1-0.2}$	$1.0^{+0.2+0.3+0.1}_{-0.2-0.2-0.2}$	$2.0^{+0.8+0.2}_{-0.6-0.2}$	$< 3.3^a$
$B^- \rightarrow b_1^0 \pi^-$	$9.6^{+0.3+1.6+2.5}_{-0.3-1.6-1.5}$	$5.1^{+3.1+3.1+0.3}_{-1.9-1.7-0.5}$	$5.0^{+1.3+0.5}_{-1.2-0.4}$	$6.7 \pm 1.7 \pm 1.0$
$\bar{B}^0 \rightarrow b_1^- \pi^+$	$0.3^{+0.1+0.1+0.3}_{-0.0-0.1-0.1}$	$1.4^{+0.4+0.1+0.1}_{-0.4-0.2-0.2}$	$0.6^{+0.3+0.1}_{-0.2-0.1}$	
$\bar{B}^0 \rightarrow b_1^+ \pi^-$	$11.2^{+0.3+2.8+2.2}_{-0.3-2.4-1.9}$	$18.7^{+9.6+8.2+1.3}_{-6.4-4.5-1.9}$	$7.7^{+2.1+0.7}_{-1.9-0.7}$	
$B^0/\bar{B}^0 \rightarrow b_1^+ \pi^-$	—	$14.8^{+8.5+6.6+1.3}_{-5.6-3.8-1.7}$	$5.0^{+1.8+0.6}_{-1.5-0.5}$	
$B^0/\bar{B}^0 \rightarrow b_1^- \pi^+$	—	$25.6^{+11.4+9.6+1.6}_{-8.3-6.0-2.6}$	$11.6^{+2.7+1.0}_{-2.5-0.9}$	
$\bar{B}^0 \rightarrow b_1^\pm \pi^\mp$	$11.4^{+0.4+2.9+2.5}_{-0.3-2.5-2.0}$	$20.2^{+9.9+8.1+1.4}_{-6.9-4.9-2.1}$	$8.3^{+2.1+0.7}_{-1.9-0.7}$	$10.9 \pm 1.2 \pm 0.9$
$\bar{B}^0 \rightarrow b_1^0 \pi^0$	$1.1^{+0.2+0.1+0.2}_{-0.2-0.1-0.2}$	$1.5^{+0.6+0.3+0.1}_{-0.5-0.3-0.2}$	$1.8^{+0.5+0.2}_{-0.4-0.1}$	$< 1.9^a$
$B^- \rightarrow b_1^0 K^-$	$6.2^{+0.5+5.0+6.4}_{-0.5-2.5-5.2}$	$24.9^{+9.8+14.9+3.7}_{-7.8-9.3-3.9}$	$4.6^{+1.9+0.7}_{-1.5-0.6}$	$9.1 \pm 1.7 \pm 1.0$
$B^- \rightarrow b_1^- \bar{K}^0$	$14.0^{+1.3+11.5+13.9}_{-1.2-5.9-8.3}$	$55.0^{+23.6+33.5+8.0}_{-17.0-21.2-8.3}$	$8.6^{+3.8+1.4}_{-3.1-1.2}$	$9.6 \pm 1.7 \pm 0.9^a$
$\bar{B}^0 \rightarrow b_1^+ K^-$	$12.1^{+1.0+9.7+12.3}_{-0.9-4.9-30.2}$	$42.9^{+17.7+26.9+6.6}_{-13.4-16.9-6.9}$	$8.5^{+3.5+1.3}_{-2.8-1.1}$	$7.4 \pm 1.0 \pm 1.0$
$\bar{B}^0 \rightarrow b_1^0 \bar{K}^0$	$7.3^{+0.5+5.4+6.7}_{-0.5-2.8-6.5}$	$23.3^{+10.6+15.5+3.5}_{-6.8-8.8-3.6}$	$4.0^{+1.8+0.7}_{-1.4-0.6}$	$< 7.8^a$

^aThe experimental data [9] is obtained on the assumption that the daughter decay $b_1 \rightarrow \pi\omega$ has a branching ratio $\mathcal{BR} = 1$.

TABLE II: Similar as table I. but for direct CP asymmetries (in %) of $B \rightarrow a_1(b_1)\pi(K)$ decays.

channel	QCDF	PQCD	SCET	Exp.
$B^- \rightarrow a_1^- \pi^0$	$0.5^{+0.3+0.6+12.0}_{-0.2-0.3-11.0}$	$1.6^{+0.0+0.1+0.2}_{-0.6-1.3-0.1}$	$-5.4^{+10.7+0.5}_{-10.1-0.5}$	$7 \pm 21 \pm 15$ $15 \pm 15 \pm 7$
$B^- \rightarrow a_1^0 \pi^-$	$-4.3^{+0.3+1.4+14.1}_{-0.3-2.2-14.5}$	$-0.9^{+0.6+0.3+0.1}_{-0.3-0.3-0.1}$	$5.7^{+11.1+0.5}_{-11.3-0.5}$	
$\bar{B}^0 \rightarrow a_1^+ \pi^-$	$-3.6^{+0.1+0.3+20.8}_{-0.1-0.5-20.2}$	$12.6^{+1.8+3.5+1.0}_{-1.2-2.5-1.1}$	$21.5^{+11.3+1.6}_{-12.7-1.9}$	
$\bar{B}^0 \rightarrow a_1^- \pi^+$	$-1.9 \pm 0.0 \pm 0.0^{+14.6}_{-14.3}$	$11.7^{+2.1+2.7+1.1}_{-1.9-2.0-1.1}$	$10.4^{+9.9+0.8}_{-10.6-0.9}$	
$\bar{B}^0 \rightarrow a_1^0 \pi^0$	$60.1^{+4.6+6.8+37.6}_{-4.9-8.3-60.7}$	$28.9^{+7.6+42.5+2.6}_{-22.1-88.1-2.5}$	$-29.5^{+15.7+2.6}_{-13.0-2.8}$	
$B^- \rightarrow a_1^- \bar{K}^0$	$0.8^{+0.0+0.1+0.6}_{-0.0-0.1-0.0}$	$-1.0^{+0.2+0.2+0.1}_{-0.0-0.2-0.1}$	$0.3^{+0.2+0.0}_{-0.2-0.0}$	$12 \pm 11 \pm 2$ $-16 \pm 12 \pm 1$
$B^- \rightarrow a_1^0 K^-$	$8.4^{+0.3+1.4+10.3}_{-0.3-1.6-12.0}$	$-6.1^{+1.1+1.3+0.6}_{-1.3-1.4-0.6}$	$-25.6^{+14.9+2.3}_{-14.8-2.4}$	
$\bar{B}^0 \rightarrow a_1^+ K^-$	$2.6^{+0.0+0.7+10.1}_{-0.1-0.7-11.0}$	$-8.9^{+1.5+2.1+0.8}_{-2.3-2.4-0.9}$	$-17.7^{+10.3+1.6}_{-10.0-1.7}$	
$\bar{B}^0 \rightarrow a_1^0 \bar{K}^0$	$-7.7^{+0.6+2.1+6.8}_{-0.6-2.2-7.0}$	$-1.8^{+0.3+0.6+0.2}_{-0.3-0.6-0.2}$	$17.9^{+10.1+1.4}_{-11.1-1.6}$	
$B^- \rightarrow b_1^- \pi^0$	$-36.5^{+4.4+18.4+82.2}_{-4.3-17.7-59.6}$	$12.5^{+16.0+23.5+1.1}_{-27.3-40.3-0.6}$	$34.1^{+27.1+3.2}_{-26.7-2.9}$	$5 \pm 16 \pm 2$
$B^- \rightarrow b_1^0 \pi^-$	$0.9^{+0.6+2.3+18.0}_{-0.4-2.7-20.5}$	$-65.8^{+11.1+13.4+4.6}_{-9.0-8.2-3.8}$	$-17.7^{+13.8+1.6}_{-14.6-1.3}$	
$\bar{B}^0 \rightarrow b_1^+ \pi^-$	$-4.0^{+0.2+0.4+26.2}_{-0.0-0.6-25.5}$	$-25.0^{+4.0+3.9+2.2}_{-4.3-4.1-1.9}$	$-42.3^{+9.3+3.6}_{-9.4-3.5}$	
$\bar{B}^0 \rightarrow b_1^- \pi^+$	$66.1^{+1.2+7.4+30.3}_{-1.4-4.8-96.6}$	$49.0^{+3.5+7.1+4.2}_{-8.0-6.0-4.0}$	0	
$\bar{B}^0 \rightarrow b_1^0 \pi^0$	$53.4^{+6.4+9.0+5.2}_{-6.3-7.3-4.7}$	$15.9^{+4.0+7.2+1.0}_{-7.8-10.7-1.4}$	$52.7^{+12.7+2.7}_{-13.9-3.9}$	
$B^- \rightarrow b_1^- \bar{K}^0$	$1.4^{+0.1+0.1+5.6}_{-0.1-0.1-0.1}$	$-0.30^{+0.02+0.00+0.03}_{-0.38-0.58-0.03}$	$-0.8^{+0.2+0.1}_{-0.2-0.1}$	$-3 \pm 15 \pm 2$ $-46 \pm 20 \pm 2$ $-7 \pm 12 \pm 2$
$B^- \rightarrow b_1^0 K^-$	$18.7^{+1.6+7.8+57.7}_{-1.7-6.1-44.9}$	$19.4^{+0.0+4.4+1.8}_{-0.4-4.0-1.8}$	$46.3^{+10.5+3.8}_{-8.8-3.9}$	
$\bar{B}^0 \rightarrow b_1^+ K^-$	$5.5^{+0.2+1.2+47.2}_{-0.3-1.2-30.2}$	$16.6^{+2.4+3.8+1.6}_{-2.3-3.2-1.5}$	$46.3^{+10.5+3.8}_{-8.8-3.9}$	
$\bar{B}^0 \rightarrow b_1^0 \bar{K}^0$	$-8.6^{+0.8+3.3+8.3}_{-0.8-4.2-25.4}$	$-4.3^{+1.6+1.8+0.4}_{-1.6-1.8-0.4}$	$-0.8^{+0.2+0.1}_{-0.2-0.1}$	

decays which can be found in the literature³. Because of the same flavor structures, the hard spectator scattering diagrams often accompany with the factorizable diagrams. One only needs to consider the flavor structure for factorizable diagrams and to use meson matrices by evaluating the master equations [38]. For the CKM matrix elements, we use the updated global fit results from CKMfitter group [35]:

$$\begin{aligned}
V_{ud} &= 0.97400, & V_{us} &= 0.22653, & |V_{ub}| &= (3.57^{+0.17}_{-0.17}) \times 10^{-3}, \\
V_{cd} &= -0.22638, & V_{cs} &= 0.97316, & V_{cb} &= (40.5^{+3.2}_{-2.9}) \times 10^{-3}, \\
\beta &= (21.7^{+0.017}_{-0.017})^\circ, & \gamma &= (67.6^{+2.8}_{-4.5})^\circ.
\end{aligned} \tag{28}$$

Predictions in the QCDF approach are also collected in the tables to make a compari-

³ There still exist two differences between the factorizable emission diagrams of $B \rightarrow AP$ and $B \rightarrow PP$ decays: the axial-vector meson can not be generated by the scalar or pseudo-scalar current, thus the chirally enhanced penguins vanish; due to the vanishing decay constant, b_1 can not be factorized from the B meson and the recoiling meson.

son [14] In the QCDF approach, a_2 (to be precise, α_2) is much smaller than 0.5, thus their amplitude from color-suppressed tree diagrams is not large enough to resolve the problem in $B^0/\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$ and $B^0 \rightarrow (a_1^- \pi^0, a_1^0 \pi^-)$ decays. Their prediction on the branching ratio of $\bar{B}^0 \rightarrow a_1^+ K^-$ is compatible with the data. For $B \rightarrow b_1 K$, they found that decay rates are sensitive to the interference between emission diagrams and annihilation diagrams. The small decay rate of $\bar{B}^0 \rightarrow b_1^+ K^-$ arises from the destructive interference between emission diagrams and annihilations, thus the prediction on branching ratio $\bar{B}^0 \rightarrow b_1^+ K^-$ is basically consistent with the data. But their predictions on four ratios of branching fractions R_{1-4} (R_3 and R_4 are related to ratios R_1 and R_2 defined in the present paper; their ratios R_1 and R_2 characterize the magnitude of color-suppressed contributions in $B \rightarrow a_1 \pi$ decay modes.) deviate from experimental data.

Several remarks on the numerical results in the PQCD approach and SCET approach are in order:

- The predictions on $\mathcal{BR}(\bar{B}^0 \rightarrow a_1^- \pi^+)$ in both approaches are a bit smaller than experimental data, because the decay constant of $f_{a_1} = 0.238$ GeV [39] is a bit smaller than that extracted from the data.
- As we expected, color-suppressed contributions to $B \rightarrow a_1 \pi$ decays are large in the SCET framework but small in the PQCD approach: SCET predictions are much larger and consistent with the present data within the uncertainties.
- In the PQCD approach, $\bar{B}^0 \rightarrow b_1^- \pi^+$ occur via the so-called hard spectator scattering diagrams, despite of the zero decay constant of b_1 . In $B^- \rightarrow b_1^0 \pi^-$, the hard spectator scattering diagrams contributions (tree operators), with a b_1^0 meson emitted, are sizable and cancel with color-allowed contribution where the pion is emitted. Thus the branching ratio of $B^- \rightarrow b_1^0 \pi^-$ is smaller than one half of $\mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ \pi^-)$.
- In the SCET approach, $\bar{B}^0 \rightarrow b_1^- \pi^+$ only receive contributions from charming penguins and correspondingly the direct CP asymmetry in this channel is 0. The predicted branching ratio is smaller than the PQCD prediction but larger than the QCDF prediction.
- In the SCET, the direct CP asymmetries in $\bar{B}^0 \rightarrow a_1^+ K^-$ and $B^- \rightarrow a_1^0 K^-$ have the same sign and similar size. Moreover, their branching ratios obey the simple relation: $\mathcal{BR}(\bar{B}^0 \rightarrow a_1^+ K^-) = 2\mathcal{BR}(B^- \rightarrow a_1^0 K^-)$. It is also similar for $\bar{B}^0 \rightarrow b_1^+ K^-$ and $B^- \rightarrow b_1^0 K^-$: the direct CP asymmetries are equal with each other; the branching ratios also satisfy the relation $\mathcal{BR}(\bar{B}^0 \rightarrow b_1^+ K^-) = 2\mathcal{BR}(B^- \rightarrow b_1^0 K^-)$, where the small deviation arises from the different mass and decay width of $\bar{B}^0$ and B^- meson.

- As expected, the two ratios R_1 and R_2 are predicted with large deviations from the data:

$$R_1 = 1.16, \quad R_2 = 0.54, \quad \text{PQCD} \quad (29)$$

$$R_1 = 0.91, \quad R_2 = 0.50. \quad \text{SCET} \quad (30)$$

- Predictions on the observables in time-dependent decay width of $B^0/\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$ and $B^0/\bar{B}^0 \rightarrow b_1^\pm \pi^\mp$ are basically consistent with the experimental data except the ΔS , the α_{eff} in $\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$ and $\mathcal{A}_{b_1\pi}$. For $\bar{B}^0 \rightarrow b_1^\pm \pi^\mp$ decays, predictions on ΔC in the two approaches are close to -1 and they are consistent with the QCDF prediction [14] and the data. In the SCET framework, the angle $\alpha_{eff}^+(\bar{B}^0 \rightarrow b_1^\pm \pi^\mp)$ is equal to $\frac{\pi}{2} - \beta$ which is also a consequence of the vanishing decay constant of b_1 meson.

VI. SUMMARY

In summary, we have investigated the $B \rightarrow a_1(b_1)\pi(K)$ decays under the factorization framework and find large differences between theoretical predictions and experimental data. In tree dominated processes $B \rightarrow a_1\pi$, large contributions from color-suppressed tree diagrams are required. In $\bar{B}^0 \rightarrow (a_1^+, b_1^+)K^-$ decays, theoretical results are larger than data by factors of 2.8 and 5.5 respectively, meanwhile ratios R_1 and R_2 defined in Eq. (19) are too much larger too. In the PQCD framework, the predicted decay rates of $B \rightarrow a_1^\pm \pi^\mp$ are consistent with data. But the other problems can not be resolved. The SCET approach has the potential to resolve the first two problems: if large hard-scattering form factors are allowed, theoretical predictions $\mathcal{BR}(B^- \rightarrow a_1^- \pi^0)$ and $\mathcal{BR}(B^- \rightarrow a_1^0 \pi^-)$ are in good agreement with data; with the help of charming penguins, large branching ratios of $\bar{B}^0 \rightarrow (a_1^+, b_1^+)K^-$ are also pulled down to the same magnitude with the data. However, the two problems on ratios in $b \rightarrow s$ transitions remain in the present theoretical methods. These two problems may indicate some new mechanism, from the non-perturbative contributions such as final state interactions or new physics scenarios, which needs further study.

Acknowledgements

This work is partly supported by National Natural Science Foundation of China under the Grant Numbers 10735080, 10625525 and 10525523. We would like to acknowledge H.Y.

TABLE III: Same as table I but for Time-dependent CP asymmetry parameters in $B^0/\bar{B}^0 \rightarrow a_1^\pm \pi^\mp$ and $B^0/\bar{B}^0 \rightarrow b_1^\pm \pi^\mp$ decays.

Observables	QCDF	PQCD	SCET	Exp.
$\mathcal{A}_{a_1\pi}$	$0.003^{+0.001+0.002+0.043}_{-0.002-0.003-0.045}$	$-0.009^{+0.002+0.002+0.001}_{-0.002-0.003-0.001}$	$0.02^{+0.08+0.00}_{-0.08-0.00}$	$-0.07 \pm 0.07 \pm 0.02$
C	$0.02^{+0.00+0.00+0.14}_{-0.00-0.00-0.14}$	$-0.12^{+0.02+0.02+0.01}_{-0.02-0.03-0.01}$	$-0.15^{+0.08+0.01}_{-0.07-0.01}$	$-0.10 \pm 0.15 \pm 0.09$
ΔC	$0.44^{+0.03+0.03+0.03}_{-0.04-0.05-0.04}$	$0.11^{+0.03+0.06+0.01}_{-0.01-0.05-0.01}$	$0.23^{+0.18+0.00}_{-0.19-0.00}$	$0.26 \pm 0.15 \pm 0.07$
S	$-0.37^{+0.01+0.05+0.09}_{-0.01-0.08-0.16}$	$-0.23^{+0.02+0.03+0.09}_{-0.01-0.03-0.14}$	$-0.45^{+0.07+0.08}_{-0.06-0.11}$	$0.37 \pm 0.21 \pm 0.07$
ΔS	$0.01^{+0.00+0.00+0.02}_{-0.00-0.00-0.02}$	$-0.03^{+0.01+0.01+0.00}_{-0.01-0.01-0.00}$	$0.02^{+0.04+0.00}_{-0.05-0.00}$	$-0.14 \pm 0.21 \pm 0.06$
α_{eff}^+	$(97.2^{+0.3+1.0+4.7}_{-0.3-0.6-2.5})^\circ$	$(93.8^{+0.4+0.7+4.4}_{-0.4-0.4-2.8})^\circ$	$(103.5^{+2.4+4.0}_{-2.5-2.6})^\circ$	
α_{eff}^-	$(107.0^{+0.5+3.6+6.6}_{-0.5-2.3-3.7})^\circ$	$(99.8^{+0.5+1.5+4.2}_{-0.7-1.4-2.7})^\circ$	$(104.8^{+2.7+4.0}_{-3.2-2.6})^\circ$	
α_{eff}	$(102.0^{+0.4+2.3+5.7}_{-0.4-1.5-3.1})^\circ$	$(96.8^{+0.4+1.0+4.3}_{-0.6-0.9-2.7})^\circ$	$(104.2^{+1.8+4.0}_{-2.0-2.6})^\circ$	$(78.6 \pm 7.3)^\circ$
$\mathcal{A}_{b_1\pi}$	$-0.06^{+0.01+0.01+0.23}_{-0.01-0.01-0.23}$	$-0.27^{+0.05+0.04+0.02}_{-0.05-0.04-0.02}$	$-0.39^{+0.08+0.03}_{-0.08-0.03}$	$-0.05 \pm 0.10 \pm 0.02$
C	$-0.03^{+0.01+0.01+0.06}_{-0.02-0.02-0.01}$	$-0.03^{+0.01+0.01+0.00}_{-0.01-0.01-0.00}$	$0.07^{+0.04+0.02}_{-0.03-0.01}$	$0.22 \pm 0.23 \pm 0.05$
ΔC	$-0.96^{+0.03+0.02+0.08}_{-0.03-0.03-0.01}$	$-0.87^{+0.02+0.04+0.01}_{-0.02-0.04-0.01}$	$-0.83^{+0.08+0.03}_{-0.07-0.03}$	$-1.04 \pm 0.23 \pm 0.08$
S	$0.05^{+0.03+0.02+0.15}_{-0.03-0.02-0.26}$	$0.08^{+0.00+0.02+0.06}_{-0.01-0.02-0.04}$	$-0.46^{+0.14+0.03}_{-0.10-0.03}$	
ΔS	$0.12^{+0.04+0.04+0.08}_{-0.03-0.04-0.09}$	$-0.24^{+0.01+0.06+0.02}_{-0.02-0.07-0.01}$	$-0.17^{+0.06+0.03}_{-0.05-0.02}$	
$\alpha_{\text{eff}}^+{}^a$	$(107.6^{+0.7+3.5+155.4}_{-0.2-4.9-17.8})^\circ$	$(174.1^{+0.0+2.5+5.0}_{-0.8-3.9-3.2})^\circ$	68.3°	
α_{eff}^-	$(101.3^{+0.4+2.1+4.9}_{-0.4-1.4-8.6})^\circ$	$(13.3^{+0.2+1.9+4.1}_{-0.3-1.7-2.7})^\circ$	$(3.4^{+0.2+4.6}_{-0.1-2.9})^\circ$	
α_{eff}	$(104.4^{+0.6+2.6+80.4}_{-0.3-2.1-1.6})^\circ$	$(93.7^{+0.0+0.9+4.5}_{-0.5-1.6-2.8})^\circ$	$(35.8^{+0.1+2.3}_{-0.0-1.5})^\circ$	

^aOne needs to be careful about the phase of the B -meson decay amplitudes [38]. For example, the $\bar{B}^0 \rightarrow b_1^- \pi^+$ and $B^0 \rightarrow b_1^+ \pi^-$ decay amplitudes are determined as:

$$A(\bar{B}^0 \rightarrow b_1^- \pi^+) = V_{ub}V_{ud}^*T + V_{cb}V_{cd}^*P, \quad A(B^0 \rightarrow b_1^+ \pi^-) = -[V_{ub}^*V_{ud}T + V_{cb}^*V_{cd}P]. \quad (31)$$

TABLE IV: Mixing-induced CP asymmetries in $\bar{B}^0 \rightarrow a_1^0 K_S$ and $\bar{B}^0 \rightarrow b_1^0 K_S$ decays.

Channel	PQCD	SCET
$\bar{B}^0 \rightarrow a_1^0 \pi^0$	$0.09^{+0.20+0.78+0.08}_{-0.19-0.87-0.08}$	$0.48^{+0.11+0.09}_{-0.14-0.15}$
$\bar{B}^0 \rightarrow a_1^0 K_S$	$0.71^{+0.01+0.02+0.00}_{-0.01-0.02-0.00}$	$0.85^{+0.05+0.01}_{-0.06-0.01}$
$\bar{B}^0 \rightarrow b_1^0 \pi^0$	$0.67^{+0.02+0.09+0.09}_{-0.00-0.06-0.07}$	$0.61^{+0.09+0.09}_{-0.11-0.06}$
$\bar{B}^0 \rightarrow b_1^0 K_S$	$-0.61^{+0.01+0.03+0.01}_{-0.01-0.03-0.01}$	-0.69

Cheng, Y.L. Shen and D.S. Yang for valuable discussions.

-
- [1] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **97**, 051802 (2006) [arXiv:hep-ex/0603050].
 - [2] K. Abe *et al.* [Belle Collaboration], arXiv:0706.3279 [hep-ex].
 - [3] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **98**, 181803 (2007) [arXiv:hep-ex/0612050].
 - [4] F. Palombo, arXiv:hep-ex/0703005.
 - [5] B. Aubert *et al.* [The BABAR Collaboration], Phys. Rev. Lett. **99**, 241803 (2007) [arXiv:0707.4561 [hep-ex]].
 - [6] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **99**, 261801 (2007) [arXiv:0708.0050 [hep-ex]].
 - [7] B. Aubert *et al.* [BABAR Collaboration], arXiv:0709.4165 [hep-ex].
 - [8] G. B. Mohanty [BABAR Collaboration], arXiv:0711.4956 [hep-ex].
 - [9] B. Aubert *et al.* [BABAR Collaboration], arXiv:0805.1217 [hep-ex].
 - [10] C. H. Chen, C. Q. Geng, Y. K. Hsiao and Z. T. Wei, Phys. Rev. D **72**, 054011 (2005) [arXiv:hep-ph/0507012].
 - [11] V. Laporta, G. Nardulli and T. N. Pham, Phys. Rev. D **74**, 054035 (2006) [Erratum-ibid. D **76**, 079903 (2007)] [arXiv:hep-ph/0602243].
 - [12] G. Calderon, J. H. Munoz and C. E. Vera, Phys. Rev. D **76**, 094019 (2007) [arXiv:0705.1181 [hep-ph]].
 - [13] K. C. Yang, Phys. Rev. D **76**, 094002 (2007) [arXiv:0705.4029 [hep-ph]].
 - [14] H. Y. Cheng and K. C. Yang, Phys. Rev. D **76**, 114020 (2007) [arXiv:0709.0137 [hep-ph]].
 - [15] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].
 - [16] E. Barberio *et al.* [Heavy Flavor Averaging Group (HFAG) Collaboration], arXiv:0704.3575 [hep-ex]. The updated results can be found at <http://www.slac.stanford.edu/xorg/hfag>.
 - [17] W. Wang, R. H. Li and C. D. Lu, arXiv:0711.0432 [hep-ph].
 - [18] K. C. Yang, unpublished; H. Hatanaka and K. C. Yang, arXiv:0804.3198 [hep-ph].
 - [19] Z. G. Wang, arXiv:0804.0907 [hep-ph].
 - [20] C. W. Bauer, D. Pirjol, I. Z. Rothstein and I. W. Stewart, Phys. Rev. D **70**, 054015 (2004) [arXiv:hep-ph/0401188].
 - [21] C. W. Bauer, I. Z. Rothstein and I. W. Stewart, Phys. Rev. D **74**, 034010 (2006) [arXiv:hep-ph/0510241].

- [22] C. W. Bauer, D. Pirjol, I. Z. Rothstein and I. W. Stewart, Phys. Rev. D **72**, 098502 (2005) [arXiv:hep-ph/0502094].
- [23] P. Colangelo, G. Nardulli, N. Paver and Riazuddin, Z. Phys. C **45**, 575 (1990).
- [24] M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. B **501** (1997) 271 [arXiv:hep-ph/9703353].
- [25] M. Ciuchini, E. Franco, G. Martinelli, M. Pierini and L. Silvestrini, Phys. Lett. B **515**, 33 (2001) [arXiv:hep-ph/0104126].
- [26] A. R. Williamson and J. Zupan, Phys. Rev. D **74**, 014003 (2006) [Erratum-ibid. D **74**, 03901 (2006)] [arXiv:hep-ph/0601214].
- [27] W. Wang, Y. M. Wang, D. S. Yang and C. D. Lu, arXiv:0801.3123 [hep-ph].
- [28] Y. Y. Keum, H. n. Li and A. I. Sanda, Phys. Lett. B **504**, 6 (2001) [arXiv:hep-ph/0004004].
- [29] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Rev. D **63**, 054008 (2001) [arXiv:hep-ph/0004173].
- [30] C. D. Lu, K. Ukai and M. Z. Yang, Phys. Rev. D **63**, 074009 (2001) [arXiv:hep-ph/0004213].
- [31] B. H. Hong and C. D. Lu, Sci. China **G49**, 357 (2006) [arXiv:hep-ph/0505020].
- [32] J. Zhu, Y.L. Shen, C.D. Lu, Phys. Rev. D **72**, 054015 (2005); H-n Li, Phys. Lett. B **622**, 68 (2005); H.W. Huang, Phys. Rev. D **73**, 014011 (2006)
- [33] H. n. Li and S. Mishima, Phys. Rev. D **73**, 114014 (2006) [arXiv:hep-ph/0602214].
- [34] M. Bona *et al.* [UTfit Collaboration], JHEP **0507**, 028 (2005) [arXiv:hep-ph/0501199].
- [35] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41**, 1 (2005) [arXiv:hep-ph/0406184].
The updated results can be found at <http://ckmfitter.in2p3.fr/>.
- [36] A. Ali, G. Kramer, Y. Li, C. D. Lu, Y. L. Shen, W. Wang and Y. M. Wang, Phys. Rev. D **76**, 074018 (2007) [arXiv:hep-ph/0703162].
- [37] R. H. Li, C. D. Lu and H. Zou, arXiv:0803.1073 [hep-ph].
- [38] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003) [arXiv:hep-ph/0308039].
- [39] K. C. Yang, Nucl. Phys. B **776**, 187 (2007) [arXiv:0705.0692 [hep-ph]].